



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $p=2rs$; $q=r^2+s^2$. Then $x=\frac{4r^2s^2}{r+s}$; $y=\frac{(r^2+s^2)^2}{r+s}$.

Let $r=k+l$; $s=k-l$. Then $x=\frac{2(k^2-l^2)^2}{k}$; $y=\frac{2(k^2+l^2)^2}{k}$.

Let $l=\alpha k$. Then $x=2k^3(1-\alpha^2)^2$; $y=2k^3(1+\alpha^2)^2$.

Now $a=p^3=8r^3s^3=8(k^2-l^2)^3=8k^6(1-\alpha^2)^3$, and $b=q^3=(r^2+s^2)^3=8(k^2+l^2)^3=8k^6(1+\alpha^2)^3$, where α and k are integers.

PROBLEMS.

53. Proposed by A. H. BELL, Box 184, Hillsboro, Illinois.

Given $x^2-114xy=\mp 3$ to find the least values of x and y in integers.

54. Proposed by JOSIAH H. DRUMMOND, LL. D., Portland, Maine.

In the expression $2x^2-2ax+b^2$, find two series of values for x in integral terms of a and b .

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

35. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

Find the chance that the distance of two points within a square shall not exceed a side of the square. [From *Byerly's Integral Calculus*.]

I. Solution by ALWYN C. SMITH, The University of Colorado, Boulder, Colorado.

a is one side of the square; P and Q the two points; (x, y) the point P with O for origin; and r and ϕ the polar coordinates of Q , with P as origin. Then the favorable cases are

$$4 \int_0^{\frac{1}{2}\pi} \int_0^a \int_0^{a-r\sin\phi} \int_0^{a-r\cos\phi} dx dy r dr d\phi = a^4(\pi - \frac{1}{6}\pi^3).$$

All the cases $= a^2 \cdot a^2 = a^4$. Therefore, $p = \pi - \frac{1}{6}\pi^3$.

